

Serial No. 10/642,438



Docket No. SACHP0145US

Claims

1. (Currently amended) An ionic liquid composition comprising at least about 55 weight percent of an ionic liquid comprising:

(a) a quatery ammonium, phosphonium or sulfonium cation having ~~more~~ than from about 4 to about 16 carbon atoms; and

(b) an anion selected from the group consisting of

**I****II**

wherein R_1 , R_2 , R_4 and R_5 are independently selected from the group consisting of substituted or unsubstituted alkyl or alkenyl groups;

wherein R_3 is a substituted or unsubstituted alkylene group, heteroarylene group, arylene group, or cycloalkylene group ;

wherein R_6 , R_7 , and R_8 are independently selected from H, alkyl, NO_2 , halo, cyano, silyl, and OH;

or R_1 and R_2 may be taken together to form a ring;

or R_4 and R_5 may be taken together to form a ring;

or R_6 and R_7 or R_7 and R_8 may be taken together to form a ring, and

Serial No. 10/642,438

Docket No. SACHP0145US

wherein when any of R₁, R₂, R₃, R₄ and R₅ are substituted, the substituent may include alkyl, NO₂, halo, cyano, silyl and OH groups.

2. (Original) The composition of Claim 1 wherein the anion has the chemical structure I.


3. (Currently amended) The composition of Claim 2 wherein R₁ and R₂ are independently selected from alkyl groups having about five or more to about eighteen carbon atoms.

4. (Cancelled)

5. (Original) The composition of Claim 2 wherein R₃ is $-(CH_2)_n-$ wherein n is an integer of from about one to about 10.

6. (Original) The composition of Claim 5 wherein R₁ and R₂ are independently selected from alkyl groups having from about six to about eighteen carbon atoms.

7. (Currently amended) The composition of Claim 6 wherein n is one and R₁ and R₂ are



8. (Original) The composition of Claim 1 wherein the anion has the chemical structure II.

9. (Original) The composition of Claim 8 wherein R₆, R₇, and R₈ are H.

Serial No. 10/642,438

Docket No. SACHP0145US

10. (Currently amended) The composition of Claim 8 wherein R_4 and R_5 are independently selected from alkyl groups having about five ~~or more~~ to about eighteen carbon atoms.

11. (Cancelled)

12. (Currently amended) The composition of Claim 9 wherein R_4 and R_5 are independently selected from alkyl groups having about five ~~or more~~ to about eighteen carbon atoms.

13. (Original) The composition of Claim 9 wherein R_4 and R_5 are $-\text{CH}_2-\text{CH}(\text{CH}_2\text{CH}_3)(\text{CH}_2-\text{CH}_2-\text{CH}_3)$.

14. (Withdrawn) The composition of Claim 2 further comprising a catalyst.

15. (Withdrawn) The composition of Claim 7 further comprising a catalyst.

16. (Original) The composition of Claim 2 further comprising a hydrocarbon.

17. (Original) The composition of Claim 7 further comprising a hydrocarbon.

18. (Withdrawn) The composition of Claim 8 further comprising a catalyst.

19. (Withdrawn) The composition of Claim 9 further comprising a catalyst.

20. (Original) The composition of Claim 8 further comprising a hydrocarbon.

Serial No. 10/642,438


Docket No. SACHP0145US

21. (Original) The composition of Claim 9 further comprising a hydrocarbon.
22. (Original) The composition of Claim 1 wherein the cation is a quaternary ammonium or quaternary phosphonium.
23. (Currently amended) The composition of Claim 22 wherein the quaternary ammonium cation is independently selected from the group consisting of substituted or unsubstituted pyridinium, pyridazinium, pyrimidinium, pyrazinium, imidazolium, pyrazolium, thiazolium, oxazolium, triazolium, imidazolinium, methylpyrrolidinium, isothiazolium, isoxazolium, oxazolium, pyrrolium, and thiophenium, wherein when substituted, the substituent group may comprise one or more of halo, alkyl, and aryl groups, and two adjacent substituents may be joined together to form an alkylene radical to form a ring structure converging on N, and in which the alkyl, phenyl and alkylene groups may be further substituted.
24. (Currently amended) The composition of Claim 1 wherein the ~~cation is an~~ quaternary ammonium cation is substituted by ~~one or more~~ groups selected from the group consisting of alkyl and aryl groups.
25. (Currently amended) The composition of Claim 22 wherein the quaternary ammonium cation is BMIM (1-butyl-3-methylimidazolium).
26. (Currently amended) The composition of Claim 1 wherein the quaternary ammonium cation is tetrabutyl ammonium, tributylmethyl ammonium, tetrabutyl phosphonium, tetraethyl ammonium, N,N-dialkyl pyrrolidinium, trimethyl-2-hydroxyethyl ammonium, N,N'-dialkyl imidazolium, N-alkylpyridinium, or a mixture thereof.
27. (Currently amended) ~~An~~ A hydrophobic ionic liquid composition comprising at least about 55 weight percent of an ionic liquid comprising:


Serial No. 10/642,438

Docket No. SACHP0145US

- (a) a quaternary ammonium, phosphonium or sulfonium cation; and
 (b) an anion selected from the group consisting of



I



II

wherein R_1 , R_2 , R_4 and R_5 are independently selected from the group consisting of substituted or unsubstituted alkyl or alkenyl groups;

wherein R_3 is a substituted or unsubstituted alkylene group, heteroarylene group, arylene group, or cycloalkylene group ;

wherein R_6 , R_7 , and R_8 are independently selected from H, alkyl, alkoxy, alkylthio, SO_3H , NO_2 , halo, cyano, silyl, and OH;

or R_1 and R_2 may be taken together to form a ring;

or R_4 and R_5 may be taken together to form a ring;

or R_6 and R_7 or R_7 and R_8 may be taken together to form a ring, and

wherein when any of R_1 , R_2 , R_3 , R_4 and R_5 are substituted, the substituent may include alkyl, NO_2 , halo, cyano, silyl and OH groups.

Serial No. 10/642,438

Docket No. SACHP0145US

28. (Cancelled)

29. (Currently amended) The composition of ~~Claim 28~~ Claim 27 wherein the cation is a quaternary ammonium or quaternary phosphonium.

30. (Currently amended) The composition of Claim 29 wherein the quaternary ammonium cation is independently selected from the group consisting of substituted or unsubstituted pyridinium, pyridazinium, pyrimidinium, pyrazinium, imidazolium, pyrazolium, thiazolium, oxazolium, triazolium, imidazolinium, methylpyrrolidinium, isothiazolium, isoxazolium, oxazolium, pyrrolium, and thiophenium, wherein when substituted, the substituent group may comprise one or more of halo, alkyl, and aryl groups, and two adjacent substituents may be joined together to form an alkylene radical to form a ring structure converging on N, and in which the alkyl, phenyl and alkylene groups may be further substituted.

31. (Currently amended) The composition of Claim 30 wherein the ~~cation is an~~ quaternary ammonium cation is substituted by ~~one or more~~ groups selected from the group consisting of alkyl and aryl groups.

32. (Currently amended) The composition of Claim 30 wherein the quaternary ammonium cation is BMIM (1-butyl-3-methylimidazolium).

33. (Currently amended) The composition of Claim 30 wherein the cation is tetrabutyl ammonium, tributylmethyl ammonium, tetrabutyl phosphonium, tetraethyl ammonium, N, N – dialkyl pyrrolidinium, trimethyl 2-hydroxyethyl ammonium, N, N' – dialkyl imidazolium, N-alkylpyridinium, or a mixture thereof.

34. (Currently amended) The composition of Claim 27 wherein the anion is Docusate (bis(2-ethylhexyl)sulfosuccinate diester).

Serial No. 10/642,438

Docket No. SACHP0145US

35. (Currently amended) The composition of Claim 27 wherein the anion has the chemical structure I ~~and is hydrophobic.~~

36. (Currently amended) The composition of Claim 27 wherein the anion has the chemical structure II ~~and is hydrophobic.~~

37. (Currently amended) The composition of ~~Claim 27~~ Claim 1 wherein the anion has the chemical structure I and is hydrophilic.

38. (Currently amended) The composition of ~~Claim 27~~ Claim 1 wherein the anion has the chemical structure II and is hydrophilic.

39. (Currently amended) The composition of Claim 1 wherein the anion has the chemical structure I and the ionic liquid is hydrophobic.

40. (Currently amended) The composition of Claim 1 wherein the anion has the chemical structure II and the ionic liquid is hydrophobic.

41. (Original) The composition of Claim 1 wherein the anion is selected from the group consisting of the anions of (i) di-n-cyclohexyl ester of sulfosuccinic acid; (ii) di-n-octyl ester of sulfosuccinic acid; (iii) di-n-butyl ester of sulfosuccinic acid; (iv) di-isobutyl ester of sulfosuccinic acid; (v) di-neopentyl ester of sulfosuccinic acid; (vi) di-n-heptyl ester of sulfosuccinic acid; and (vii) di-n-heptyl ester of sulfosuccinic acid.

42. (Original) The composition of Claim 41 wherein the cation is tetrabutyl ammonium.

Serial No. 10/642,438

Docket No. SACHP0145US

43. An ionic liquid composition, comprising at least about 55 weight percent of an ionic liquid comprising:

- (a) a quaternary ammonium, phosphonium or sulfonium an-onium cation; and
(b) an anion having the following structure:



III

wherein R_1 , R_2 , R_3 , R_4 , and R_5 are independently selected from the group consisting of a hydrogen atom and a carbon-containing group;
and wherein the ionic liquid has a melting point that is less than about 100°C.

44. (Original) The composition of Claim 43, wherein R_1 is 2-ethylhexyl, R_2 is ethyl, R_3 is a methylene group, R_4 is ethyl, and R_5 is 2-ethylhexyl.

45. (Original) The composition of Claim 44, wherein the cation is tetrabutyl ammonium.

46. (Original) The composition of Claim 44, wherein the cation is 1-methyl-3-hexyl imidazolium.

47. (Original) The composition of Claim 43, wherein R_1 is 2-ethylhexyl, R_2 is a hydrogen atom, R_3 is a methylene group, R_4 is a hydrogen atom, and R_5 is 2-ethylhexyl.

48. (Original) The composition of Claim 47, wherein the cation is tetrabutyl ammonium.

49. (Original) The composition of Claim 47, wherein the cation is 1-methyl-3-hexyl imidazolium.

50. (Original) The composition of claim 43, further comprising a hydrocarbon.

Serial No. 10/642,438

Docket No. SACHP0145US

51. (Currently amended) The composition of claim 1, wherein the cation and the anion form a molten salt having a melting point of less than about 100°C, the molten salt being selected from the group consisting of tetrabutylammonium docusate, MeBu₃N docusate, Me₃N(CH₂)₆NMe₃ docusate, Bu₄P docusate, Et₄N docusate, 1-hexyl-3-methyl imidazolium docusate, 1-octyl-3-methyl imidazolium bromide docusate, 1-butyl-3-methyl imidazolium docusate, and 1-methyl-2-ethyl imidazolium docusate, wherein docusate is bis(2-ethylhexyl)sulfosuccinate diester.

52. (Currently amended) A composition comprising:
(a) an ionic liquid ~~containing~~ comprising an anion that is selected from the group consisting of (i) docusate, (ii) an anion of a bis(organo)ester derivative of sulfosuccinic acid, and (iii) an anion of a bis(organoamide) derivative of sulfosuccinic acid and a quaternary ammonium, phosphonium or sulfonium cation; and
(b) CO₂ at supercritical conditions;
wherein the ionic liquid is dissolved in the CO₂.

53. (Withdrawn) A composition comprising:
(a) a hydrocarbon fuel; and
(b) an ionic liquid containing an anion that is selected from the group consisting of (i) docusate, (ii) an anion of a bis(organo)ester derivative of sulfosuccinic acid, and (iii) an anion of a bis(organoamide) derivative of sulfosuccinic acid.

54. (Withdrawn) A composition comprising:
(a) a polymer; and
(b) an anti-static additive comprising an ionic liquid containing an anion that is selected from the group consisting of (i) docusate, (ii) an anion of a bis(organo)ester derivative of sulfosuccinic acid, and (iii) an anion of a bis(organoamide) derivative of sulfosuccinic acid.

Serial No. 10/642,438

Docket No. SACHP0145US

55. (Withdrawn) The composition of claim 54, wherein the polymer is polyvinylacetate.

56. (Cancelled)

57. (Currently amended) The ionic liquid composition of ~~claim 56~~ claim 1 wherein the ionic liquid melts at a temperature range that is greater than about 40° C but less than about 80° C.

58. (Currently amended) A composition, comprising:
a first ionic liquid combined with a second ionic liquid,

(a) the first ionic liquid comprising:

(i) a cation selected from the group consisting of quaternary ammonium, sulfonium, and phosphonium cations, said cation being non-tetrahedrally symmetric;

(ii) an anion having the formula Al_yR_{3y+1} wherein y is greater than 0 and R is independently selected from the group consisting of an alkyl group and halogen group;

(b) the second ionic liquid comprising an anion that is selected from the group consisting of (i) docusate, (ii) an anion of a bis(organo)ester derivative of sulfosuccinic acid, and (iii) an anion of a bis(organoamide) derivative of sulfosuccinic acid.

59. (Original) The composition of claim 58, further comprising reactants, the first and second ionic liquids being an effective reaction solvent for the reactants.